arXiv:0708.3443v2 [math.MG] 22 Nov 2007

THE SPECIAL CUTS OF THE 600-CELL

MATHIEU DUTOUR SIKIRIĆ AND WENDY MYRVOLD

Abstract. A polytope is called *regular-faced* if every one of its facets is a regular polytope. The 4-dimensional regular-faced polytopes were determined by G. Blind and R. Blind [2, 3, 4]. The last class of such polytopes is the one which consists of polytopes obtained by removing a set of non-adjacent vertices (an independent set) of the 600-cell. These independent sets are enumerated up to isomorphism and it is determined that the number of polytopes in this last class is $314,248,344$.

1. Introduction

A *d-dimensional polytope* is the convex hull of a finite number of vertices in $\mathbb{R}^d$. A d-dimensional polytope is called *regular* if its isometry group is transitive on flags. The regular polytopes are (see, for example, [8]) the regular n-gon, d-dimensional simplex α_d, hypercube γ_d, cross-polytope β_d, the 3-dimensional Icosahedron Ico, Dodecahedron Dod, the 4-dimensional 600-cell, 120-cell and 24-cell.

A polytope is called *regular-faced* if its facets are regular polytopes. If, in addition, its symmetry group is vertex-transitive then it is called *semiregular*. Several authors have considered this geometric generalization of the regular polytopes. An overview of this topic has been given by Martini [13, 14]. The 3-dimensional regular-faced polytopes have been determined by Johnson [11] and Zalgaller [19]; see [1, 10, 18, 15] for some beautiful presentation. Three papers [3, 4, 2] give a complete enumeration for the cases with dimension $d \geq 4$. G. Blind and R. Blind [6] characterized the semiregular polytopes.

Given a d-dimensional regular polytope P, $Pyr(P)$ denotes, if it exists, the regular faced $(d+1)$-dimensional polytope obtained by taking the convex hull of P and a special vertex v. The *bipyramid* $BPyr(P)$ denotes a $(d+1)$-dimensional polytope defined as the convex hull of P and two vertices, v_1 and v_2 on each side of P. The list of regular-faced d-polytopes for $d \geq 4$ is:

1. the regular d-polytopes,
2. two infinite families of d-polytopes ($Pyr(\beta_{d-1})$ and $BPyr(\alpha_{d-1})$),

(3) the semiregular polytopes n_{21} with $n \in \{0, 1, 2, 3, 4\}$ of dimension $n + 4$ and the semiregular 4-dimensional octicosahedric polytope,
(4) three 4-polytopes ($Pyr(Ico)$, $BPyr(Ico)$ and the union of $0_{21} + Pyr(\beta_3)$, where β_3 is a facet of 0_{21}), and
(5) any *special cut* 4-polytope, arising from the 600-cell by the following procedure: if C is a subset of the 120 vertices of the 600-cell, such that any two vertices in C are not adjacent, then the special cut 600_C is the convex hull of all vertices of the 600-cell, except those in C.

This paper describes the enumeration of all such special cuts (see Table 1 and [9] for the results). The enumeration of special cuts with 2, 23 and 24 vertices is done in [5]. The ones with 3, 4, 5, 6, 21, and 22 vertices are enumerated by Kirrmann [12]. Also, Martini [13] enumerated the number of special cuts with n vertices for $n \leq 6$.

2. GEOMETRY OF SPECIAL CUTS

The 600-cell has 120 vertices, and its symmetry group is the Coxeter group H_4 with $14,400$ elements. A subset C of the vertex set of the 600-cell is called *independent* if any two vertices in C are not adjacent. Given an independent subset C of the vertex-set of 600-cell, denote by 600_C the polytope obtained by taking the convex hull of the remaining vertices. Two polytopes 600_C and $600_{C'}$ are isomorphic if and only if C and C' are equivalent under H_4.

If C is reduced to a vertex v, then the 20 3-dimensional simplex facets containing v are transformed into an icosahedral facet of $600_{\{v\}}$, which we denote by Ico_v. It is easy to see that if one takes two vertices v and v' of the 600-cell, then the set of simplices containing v and v' are disjoint if and only if v and v' are not adjacent. Therefore, if C is an independent set of the vertices of the 600-cell then 600_C is regular-faced and is called a *special cut.* The name special cut come from the fact that 600_C can be obtained from the 600-cell by cutting it with the hyperplanes corresponding to the facet defined by the icosahedra Ico_v for $v \in C$.

A vertex v of a special cut 600_C can be contained in at most 3 icosahedra $(Ico_w)_{w \in C}$. The possible ways of having a vertex of 600_C contained in an icosahedron Ico_w are listed in Figure 1. It is easy to see that an independent set C has at most 24 vertices. The number of special cuts on up to 24 vertices is listed in Table 1 organized in column of cliques with the same order of symmetry group. A special cut 600_C is called *maximal* if we cannot add any vertices to C and still have a

	1	2	3	4	5	6	8	9	10	12	16	18	20
1													
2							1			2			3
3	1	21		6		3	1		1	2			3
4	187	184	2	40		7	6			3			2
5	3721	938	4	79		21	3		1	7			1
6	41551	3924	17	212		34	18		6	8			
7	321809	12093	53	322		63	4		19	12			4
8	1792727	32714	102	672	1	102	40		28	17	3		
9	7284325	70006	170	815		137	6		14	19		1	2
10	21539704	129924	282	1349	2	190	43		4	16			3
11	45979736	194232	420	1346		251	6		11	15			3
12	69895468	247136	505	1781		236	57	1	37	21	4	1	12
13	74365276	252040	527	1457		266	6		58	20			7
14	54266201	213377	553	1545		255	43		26	31			9
15	26605433	142212	478	1041	2	181	4	1	5	19		1	4
16	8612476	76249	316	837		165	39		5	14	4		
17	1824397	31465	216	461		116	4		16	6			3
18	252764	10001	123	273		45	20		25	10		1	
19	22673	2360	49	120		39	3		12	8			1
20	1202	388	18	40		17	5		1	7			
21	22	37	6	12		5	1					1	
22						5	1						
23													
24													

	24	30	32	36	40	48	72	100	120	144	192	240	576
1									1				
2												1	
3				1									
4	3		1		1								
5								1					
6	2			1		1	1						
7	1												
8	6					2					1		
9	2			1									
10	8				1			1					
11													
12	5				1	2			1	1			
13									2				
14	7				1				1				
15	2	1											
16	4					1					1		
17	1												
18	4			1		1							
19													
20	2		1			1						1	
21	2												
22	2					1							
23	1												
24													1

Table 1. The number of special cuts between 1 and 24 vertices with the orders of their symmetry groups

	1	2	3	4	5	6	8	10	12	16
10										
11		1								
12	18	9		4			1			1
13	1555	146		23						
14	39597	980		52		4	4			
15	221823	2997	9	64	2	4		3	1	
16	341592	4573	10	113		16	7		11	1
17	192266	4081	9	59		7				
18	49741	2251	19	54		26	8		2	
19	6771	838	7	39		7		6		
20	598	199	6	14		12	2	1	5	
21	17	20	2	11						
22						3				
24										

	18	20	24	30	40	48	100	144	240	576
10							1			
11										
12			1					1		
13										
14		2			1					
15		1		1						
16						1				
17										
18	1		1							
19										
20			1			1			1	
21										
22			2							
24										1

TABLE 2. The number of maximal special cuts between 10 and 24 vertices with the orders of their symmetry groups


Case I Case II Case III Case IV Case V

FIGURE 1. The local possibilities for vertices

special cut; a list of these is given in Table 2 again with the symmetry group informations.

Table 3 provides some information about highly symmetric special cuts. The column "conn." refers to the connectivity of the graph defined by the simplices of 600_C with two simplices adjacent if they

$\|V\|$	$\|G\|$	maximal	conn.	vertex orbits
24	576	yes	no	$(96, V, C_{3v})$
20	240	yes	no	$(40, V, C_{3v})$, $(60, IV, C_{2v})$
16	192	no	yes	$(96, IV, C_s)$, $(8, I, T_h)$
8	192	no	yes	$(96, II, C_s)$, $(16, I, T)$
12	144	yes	yes	$(36, IV, C_{2v})$, $(72, II, C_s)$
10	100	yes	yes	$(100, II, C_1)$, $(10, III, D_5)$

Table 3. Some highly symmetric special cuts

share a 2-dimensional face. In the column "vertex orbits", the sizes of the vertex orbits, their types according to Figure 1 and the nature of the vertex stabilizer according to its Schoenflies symbol are listed. The 143 cases with at least 20 symmetries are available from [9].

- The Snub 24-cell is the semiregular polytope obtained as 600_C with $|C| = 24$. Its symmetry group has order 576 and its facets are 24 icosahedra and 120 3-dimensional simplices in two orbits O_1, O_2 with $|O_1| = 24$, and $|O_2| = 96$. The simplices in O_1 are adjacent only to simplices in O_2. The 24 vertices of C form a 24-cell, and hence the name *snub 24-cell*. Coxeter [8] provides further details.
- The vertex set of the 24-cell can be split into three cross-polytopes β_4. Selecting one or two of these cross-polytopes gives two special cuts with 8 and 16 vertices and 192 symmetries.
- It is easy to see that the minimum size of a maximal special cut is at least 10. One of size 10 can be constructed as follows (indicating that the minimum size of a maximal cut is 10). The vertex set of the 600-cell is partitioned into two cycles of 10 vertices each and a set containing the 100-remaining vertices. The convex hull of the 100 remaining vertices is called a *Grand Antiprism* (discovered by Conway [7]). Taking a maximum independent set of each of these cycles (a total of 10 vertices since five are selected from each cycle) gives the unique (up to isomorphism) maximal special cut of order 10.

3. Enumeration methods

The *skeleton* of a polytope P is the graph formed by its vertices and edges. Enumerating the special cuts of the 600-cell is the same as enumerating the independent sets of its skeleton.

In order to ensure correctness, the independent sets were enumerated by two entirely different methods and the results were checked to ensure that they agreed. The first method used to enumerate the independent sets was a parent-child search (see [16]). The 120 vertices of 600-cell were numbered and then the search considered only the independent sets which were lexicographically minimum in their orbit. The method is then the following: given a lexicographically minimum independent set S, we consider all ways to add a vertex v such that $v > \max(S)$ and $S \cup \{v\}$ is still a lexicographically minimum independent set. Given a lexicographically minimum independent set $S = \{v_1, v_2, \dots, v_k\}$ with $v_i < v_{i+1}$, this method provides a canonical path to obtain S; first obtain $\{v_1\}$, then $\{v_2\}$, until one gets S.

The second method is explained in [17]. The algorithm uses a novel algorithmic trick combined with appropriate data structures to decrease the running time of the search. One advantageous feature of this algorithm is that the symmetries of the independent sets generated are available with no additional computation required. This method proved much faster by a factor of 1000; the relationship between the performance difference which can be attributed to the algorithm versus the quality of the programming has not been determined.

References

[1] M. Berman, *Regular-faced convex polyhedra*, J. Franklin Inst. **291** (1971) 329–352.

[2] G. Blind and R. Blind, *Die Konvexen Polytope im* $\mathbb{R}^4$*, bei denen alle Facetten reguläre Tetraeder sind*, Monatshefte für Mathematik **89** (1980) 87–93.

[3] R. Blind, *Konvexe Polytope mit regulären Facetten im* R^n $(n \geq 4)$., Contributions to geometry (Proc. Geom. Sympos., Siegen, 1978), pp. 248–254, Birkhäuser, Basel-Boston, Mass., 1979.

[4] R. Blind, *Konvexe Polytope mit kongruenten regulären* $(n-1)$*-Seiten im* R^n $(n \geq 4)$, Comment. Math. Helv. **54-2** (1979) 304–308.

[5] G. Blind and R. Blind, *Über die Symmetriegruppen von regulärseitigen Polytopen*, Mh. Math. **108** (1989) 103–114.

[6] G. Blind and R. Blind, *The semiregular polytopes*, Commentari Mathematici Helvetici **66** (1991) 150–154.

[7] J.H. Conway, *Four-dimensional Archimedean polytopes*, Proc. Colloquium on Convexity, Copenhagen 1965, Kobenhavns Univ. Mat. Institut (1967) 38–39.

[8] H.S.M. Coxeter, *Regular Polytopes*, 3rd ed., Dover, New York, 1973.

[9] M. Dutour, http://www.liga.ens.fr/~dutour/SpecialCuts/

[10] S. Gagnon, *Convex polyhedra with regular faces*, Structural Topology **6** (1982) 83–95.

[11] N.W. Johnson, *Convex polyhedra with regular faces*, Canadian Journal of Mathematics **18** (1966) 169–200.

[12] G. Kirrmann, *Regulärseitige Polytope, die durch Abschneiden von Ecken eines 600-Zells entstehen*, University of Stuttgart, 1992.

[13] H. Martini, *A hierarchical classification of Euclidean polytopes with regularity properties. Polytopes: abstract, convex and computational*, (Scarborough, ON,

1993), 71–96, NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., 440, Kluwer Acad. Publ., Dordrecht, 1994.

[14] H. Martini, *Reguläre Polytope und Verallgemeinerungen*, in: Geometrie und ihre Anwendungen. Hrsg. O Giering ung J. Hoschek, Carl Hanser Verlag, München, 1994, 247–281.

[15] Wolfram Inc., http://mathworld.wolfram.com/JohnsonSolid.html

[16] B.D. McKay, *Isomorph-free exhaustive generation*, J. Algorithms **26-2** (1998) 306–324.

[17] W. Myrvold and P.W. Fowler, *Fast enumeration of all independent sets of a graph*, preprint, (2007).

[18] A. Pugh, *Polyhedra: A Visual Approach*, University of California Press, Berkeley, 1976.

[19] V. Zalgaller, *Convex polyhedra with regular faces*, Translated from Russian. Seminars in Mathematics, V. A. Steklov Mathematical Institute, Leningrad, Vol. 2 Consultants Bureau, New York 1969 vi+95 pp.

Mathieu Dutour Sikirić, Rudjer Bosković Institute, Bijenicka 54, 10000 Zagreb, Croatia
E-mail address: `Mathieu.Dutour@ens.fr`

Wendy Myrvold, Dept of Computer Science, University of Victoria, P.O. Box 3055, Stn CSC, Victoria, B.C. Canada V8W 3P6
E-mail address: `wendym@cs.uvic.ca`